

# Kawasaki Disease

## What is Kawasaki Disease?

and spinal cord. Vasculitis causes inflammation of the blood vessels. This restricts blood flow and can damage the brain and

spinal cord, and can cause loss of brain function, stroke and even life-threatening conditions.

Vasculitis is classified as an autoimmune disorder, which occurs when the body's natural defense system mistakenly attacks

healthy tissues.

CNSV is typically categorized as primary and secondary:

Primary CNSV is confined specifically to the brain and spinal cord. It is not associated with any other systemic disease

(affecting the whole body).

Secondary CNSV usually occurs in the presence of other autoimmune diseases such as systemic lupus erythematosus or

Sjögren's syndrome and systemic forms of vasculitis such as granulomatosis with polyangiitis, microscopic polyangiitis,

Behçet's syndrome or viral or bacterial infections.

## Causes

*The cause of CNSV is not fully understood. It is believed that several factors may be involved:*

*An infection may contribute to the onset of CNSV.*

*Environmental and genetic factors may play a role as well.*

## Who Gets CNSV?

*In general, CNSV is considered rare. In the case of primary CNSV, the disorder can affect people of all ages but generally peaks around age 50. It most often occurs in males.*

## Symptoms

- Many forms of vasculitis are accompanied by fever,
- fatigue, and unintentional weight loss. CNSV symptoms
- may also include:
- Chronic headaches
- Strokes or transient ischemic attacks (mini-strokes)
- Swelling of the brain (encephalopathy)
- Forgetfulness or confusion
- Muscle weakness or paralysis
- Difficulty with coordination
- Abnormal sensations or loss of sensations
- Vision problems
- Seizures or convulsions

## Complications

The most serious complication is heart involvement caused by inflammation of the coronary arteries (vessels that bring blood to the heart muscle). This can result in aneurysms (an abnormal bulging of the arteries that can rupture). Other potential complications include inflammation of the heart muscle, heart valve problems or, very rarely, heart attack.

## Diagnosis

There is no single test for diagnosing Kawasaki disease, so your child's doctor will consider several factors including a detailed medical history, physical examination, laboratory tests, specialized imaging studies, and, when indicated, a biopsy of an affected tissue or organ.

The presence of classic symptoms helps point to a diagnosis of Kawasaki disease including fever lasting at least five days, and at least four of five of the following:

- Swollen lymph nodes
- Bloodshot eyes
- Swollen lips/mouth/tongue
- Swollen hands and feet
- Rash

The doctor will try to rule out other causes of fever or diseases with similar symptoms such as measles, scarlet fever, Rocky Mountain spotted fever, and juvenile rheumatoid arthritis.

If Kawasaki disease is suspected, your child's doctor will likely order blood and urine tests, as well as imaging tests to look for changes or damage to the heart. Common diagnostic tests to detect potential heart involvement include the echocardiogram (a cardiac ultrasound), which checks the heart valves and function, and electrocardiogram (EKG), which records the electrical activities of heart muscle. Echocardiograms or other diagnostic tests may need to be repeated at certain intervals to monitor for heart damage following recovery from this disease.

## Treatment

To prevent complications, treatment should begin as soon as possible after diagnosis, ideally within the first 10 days. Most children with Kawasaki disease require treatment in a hospital. Standard treatment includes high doses of IVIG, a human blood protein that helps prevent damage to the heart blood vessels, if given early in the disease. Aspirin is given to reduce fever, rash, joint swelling, and pain. (Note: Never give aspirin to your child without a doctor's orders. Aspirin use in children and teenagers has been associated with Reye's syndrome, a rare but serious condition that causes swelling in the brain and liver.)

Patients usually improve significantly within 24 hours of treatment with IVIG. However, some patients may need to take corticosteroids or other immunosuppressant medications if initial treatments aren't working. Rarely, surgery may be required to repair damaged heart blood vessels.

## Side Effects of Treatment

The medications used to treat Kawasaki disease have potentially serious side effects, so it is important to monitor your child's symptoms and report them to the doctor. Infection prevention is also very important. Talk to the doctor about ageappropriate vaccinations, which can reduce your child's risk of infection.

## Medical Follow-up/Relapse

Even with effective treatment, relapses can occur with Kawasaki disease. If your child's fever or other symptoms return, report them to the doctor as soon as possible. Regular checkups and ongoing monitoring of laboratory and imaging tests are important in detecting relapses or new organ involvement.

## Your Child's Medical Team

Effective treatment of Kawasaki disease may require the coordinated efforts and ongoing care of a team of medical providers and specialists. In addition to a pediatrician or primary care doctor, your child may need to see the following specialists:

- Pediatric cardiologist (specialist in diagnosis and treatment of heart disorders in children)
- Pediatric or adult infectious disease specialist (to help rule out infectious disease as a cause of fever)
- Pediatric or adult rheumatologist (joints and immune system)
- Pediatric dermatologist (skin) or others as needed

The best way to manage Kawasaki disease is to actively partner with your child's health care providers. Get to know the members of the health care team. It may be helpful to keep a health care journal to track your child's medications, symptoms, test results, and notes from doctor appointments in one place. To get the most out of doctor visits, make a list of questions beforehand and bring along a supportive friend or family member if necessary to provide a second set of ears and take notes.

Remember, it's up to you to be your child's advocate. If you have concerns about the treatment plan, be sure to speak with the medical team. It's always your right to seek a second opinion.

## Outlook

Most children with Kawasaki disease experience a full recovery. However, one in four may develop serious heart problems, so early diagnosis and treatment are critical in preventing heart damage. A very small percentage of patients die of complications from coronary blood vessel inflammation.

Children with Kawasaki disease should have follow-up echocardiograms and checkups to screen for ongoing heart problems.

In 2021 the American College of Rheumatology (ACR) published guidelines for the management of certain vasculitides, that were also endorsed by the Vasculitis Foundation (VF). Clinical practice guidelines are developed to reduce inappropriate care, minimize geographic variations in practice patterns, and enable effective use of health care resources. Guidelines and recommendations developed and/or endorsed by the ACR are intended to provide guidance for particular patterns of practice and not to dictate the care of a particular patient. The application of these guidelines should be made by the physician in light of each patient's individual circumstances. Guidelines and recommendations are subject to periodic revision as warranted by the evolution of medical knowledge, technology, and practice.

## About Vasculitis

Vasculitis is a family of nearly 20 rare diseases characterized by inflammation of the blood vessels, which can restrict blood flow and damage vital organs and tissues. Vasculitis is classified as an autoimmune disorder, which occurs when the body's natural defense system mistakenly attacks healthy tissues. Triggers may include infection, medication, genetic or environmental factors, allergic reactions, or another disease. However, the exact cause is often unknown.

## A Family of Diseases

- Anti-GBM disease (formerly Goodpasture's syndrome)
- Aortitis
- Behçet's syndrome
- Central nervous system vasculitis (CNSV)
- Cogan's syndrome
- Cryoglobulinemic vasculitis
- Cutaneous small-vessel vasculitis (CSVV) (formerly hypersensitivity/leukocytoclastic)
- Eosinophilic granulomatosis with polyangiitis (EGPA, formerly Churg-Strauss syndrome)
- Giant cell arteritis (GCA)
- Granulomatosis with polyangiitis (GPA, formerly Wegener's)
- IgA vasculitis (formerly Henoch-Schönlein purpura)
- Kawasaki disease
- Microscopic polyangiitis (MPA)
- Polyarteritis nodosa (PAN)
- Polymyalgia rheumatica (PMR)
- Rheumatoid vasculitis
- Takayasu arteritis (TAK)
- Urticarial vasculitis
- (normocomplementemic or hypocomplementemic)

## About the VF

The VF is the leading organization in the world dedicated to diagnosing, treating, and curing all forms of vasculitis. The VF is a 501(c)(3) nonprofit organization governed by a Board of Directors and advised on medical issues by a Medical and Scientific Advisory Board. VF's educational materials are not intended to replace the counsel of a physician. The VF does not endorse any medications, products, or treatments for vasculitis, and advises you to consult a physician before initiating any treatment.

The VF gratefully acknowledges Alexandra Villa-Forte, MD, Cleveland Clinic Center for Vasculitis Care and Research, for her expertise and contribution to this brochure.

To access additional VF support and educational resources, please scan the QR code below.

## VF Mission

Building upon the collective strength of the vasculitis community, the Foundation supports, inspires and empowers individuals with vasculitis, and their families, through a wide range of education, research, clinical, and awareness initiatives.



PO Box 28660, Kansas City, Missouri 64188-8660 • Phone: 816.436.8211 • Toll Free: 800.277.9474 Email:  
**[vf@vasculitisfoundation.org](mailto:vf@vasculitisfoundation.org) • [www.VasculitisFoundation.org](http://www.VasculitisFoundation.org)**

©2023 • Reproduction of this material is by permission only. • Revised September 2023